

## Neck pain and hand grip strength in dentists of Saudi Arabia

*Eman Samir Fayez*

Assistant professor in department of Physical Therapy,  
Faculty of Applied Medical Science, University of Dammam, KSA

Copyright © 2014 ISSR Journals. This is an open access article distributed under the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**ABSTRACT:** Hand grip is the functional unit of dentists used in firm holding of instruments and tools. Muscles of hand grip supplied by cervical nerves. Dentists put their neck in high physical load for prolonged time that lead to neck pain. 54.4% of dentists in K.S.A. have neck pain and 21% drop out of their job. *Objective:* The main objective is to correlate neck pain intensity with hand grip strength in dentists of Saudi Arabia. *Methodology:* *Subjects:* 25 chronic neck pain dentists (suffering from neck pain more than 3 months) in both gender, were selected. Their ages ranged from 25 to 40 years with normal BMI (18.5-25).they were divided into 3 subgroups according to neck position during their work. *Procedures:* Pain intensity will be determined by visual analog scale (VAS), hand grip strength will be measured by Jamar hand dynamometer in kilogram (Kg) unit. *Result:* Among 25 dentists the result showed extremely significant correlation between neck pain intensity and hand grip strength (direct proportional). *Discussion and conclusion:* the findings of this study denote presence of direct proportional correlation between neck pain intensity and hand grip strength which may be attributed to deficits in sensory motor integration and neuro hyper excitability found in neck pain victims.

**KEYWORDS:** neck pain, dentist, hand grip strength, sensory motor integration.

### 1 INTRODUCTION

Neck pain is an ache or discomfort in anatomical area between occiput and 3<sup>rd</sup> thoracic vertebra and laterally between middle margin of scapula<sup>(1)</sup>. Neck pain is considered to be chronic if the patient suffered more than 3 months<sup>(2, 3)</sup>. The prevalence for neck pain varies between 16.7% and 75.1% in the general population<sup>(4)</sup>. Neck pain may be caused by; spondylosis, spinal stenosis, disc herniation, stress, poor posture, prolonged posture<sup>(5, 6)</sup>. Symptoms of neck pain may include; neck soreness and headache, pain around shoulder blades, arm complaints (pain, numbness or weakness).

Dental profession is one of the occupational risk factors of neck pain<sup>(1, 7)</sup>. Studies showed that neck pain starts early in dental career even during educational training<sup>(8, 9)</sup>. In a recent study, 54.4% of dentists in K.S.A. had neck pain, 21.69% of them drop out of job<sup>(10)</sup>.

Recently, it was shown that there is a strong relation between work duration and neck pain in the staff of dental school<sup>(11, 12)</sup>. Another study reported that increasing weekly work hours has little effect on neck pain<sup>(10, 13)</sup>. The most common cause of neck pain in dentists is prolonged work posture which makes high load on muscles of neck and shoulder region (trapezius more than splenius)<sup>(12)</sup>. The main working posture of dentist is in sitting, with neck flexion, shoulder elevation and upper extremity abduction and his patient supine<sup>(10, 11, 13, 14)</sup>.

Hand is the functional unit of the dentist and is used in keeping firm holding of instruments and tools<sup>(15)</sup>. Hand grip is the amount of static force that the hand can squeeze around dynamometer or an index of the power which the hand can exert<sup>(16, 17)</sup>. Grip strength can be affected by gender (male > female), age (peaked at 4<sup>th</sup> decade), dominant more than non-dominant hand<sup>(16-20)</sup>. Different positions may also affect hand grip strength: elbow during measurement (extension > flexion 90%), supported forearm more than unsupported and standing position more than sitting<sup>(15, 21)</sup>. Grip strength used to assess functional activity of hand<sup>(22)</sup>. The measurement of hand grip strength used to detect whole upper arm strength<sup>(23)</sup>. Studies

show that persons with unilateral upper limb chronic pain have strength reduction of hand grip 20-30% less than non-painful side. In addition to that, unilateral musculoskeletal pain cause delay grip initiation and relaxation<sup>(24, 25)</sup>.

From these literature reviews we conclude that neck pain is in high prevalence in dentists and most of them take neck flexion during their work which leads to compression of nerves supply upper limb. The hand grip is a functional unit of their carrier, No previous studies show the correlation between neck pain and hand grip of dentists. So, the purpose of our study is to correlate between neck pain and hand grip strength in dentists of K.S.A.

### 1.1 THE PURPOSE OF THE STUDY

The purposes of the present study were to determine the relationship between neck pain intensity and hand grip strength in dentists of Saudi Arabia.

## 2 PROCEDURES

### 2.1 SUBJECTS SELECTION

25 chronic neck pain dentists in both gender ( 13 male and 12 female) were recruited from dental clinic of university of Dammam and several private clinics.

### 2.2 INCLUSION CRITERIA

Twenty five dentists suffered from neck pain more than three months. Both genders. Their age range from 25 to 40 years with mean age  $31.32 \pm 4.87$  years. Normal Body mass index ( $18.5-25 \text{ Kg/cm}^2$ )

### 2.3 EXCLUSION CRITERIA


The participants were excluded if they had a history of either orthopedic or neurologic diseases causing functional defect of the hand strength like: fracture, surgery of upper limb or hand, carpal tunnel syndrome, De Quervain's syndrome or diabetes mellitus.

### 2.4 INSTRUMENTS

1. Data collection sheet which include (personal data, main neck position during working). It was used for gathering data from dentists.
2. Visual analog scale (VAS) to measure the intensity of pain<sup>(26, 27)</sup>.
- 3- Jamar hand dynamometer to measure hand grip strength<sup>(16,17,19)</sup>.



*Figure 1*



*Figure 2*

## 2.5 PRE ASSESSMENT INSTRUCTION

Cease taking analgesic drugs 2-3 days before examination.

## 2.6 STEPS OF ASSESSMENT

1-Intensity of pain was determined by VAS rating from (0= no pain, 10= unbearable pain).

2-The main neck position mainly used by each dentist during work was determined and mentioned in the data collection sheet.

3-Grip strength for each dentist was measured in sitting position on chair with back support and armrest, shoulder adducted, elbow 90 degree flexed, forearm and wrist in mid position. The participant held the dynamometer by dominant hand ( figure 2). Therapist gave the command "squeeze around dynamometer as much as you can". Therapist was stand in front of dentist to read the amount of force. The mean of three measurements was taken in Kg unit, with one minute rest between each squeeze to overcome fatigue<sup>(15, 16, 20)</sup>.

## 2.7 DATA ANALYSIS

1. Data was collected from the sheet and the result of assessment and entered to SPSS program.
2. Descriptive analysis (mean, standard deviation and P value).
3. Regression analysis to find the correlation between intensity of neck pain and strength of hand grip.
4. One way ANOVA to compare the intensity of pain and grip strength in different neck position.

## 3 RESULT

The results of this study showed extremely significant correlation between neck pain and hand grip strength ( direct proportional )with p-value ( 0.0001)and the slope is significantly different from zero as seen in table" 1" and figure" 3". The mean of grip strength is  $(33.42 \pm 13.05)$  kg. The mean of pain intensity is  $(3.58 \pm 1.88)$ .

Table" 2" and figure "4" showed that there was no significant difference between pain intensity in different neck positions during working of dentists by using one-way ANOVA with p-value (0.662), but neck rotation combined with flexion took the highest pain intensity  $(4.33 \pm 3.3)$ .

Table "3" and figure " 5 " showed that there was no significant difference between hand grip strength in different neck positions by using one –way ANOVA with p-value (0.196). Neck rotation also took highest hand grip strength  $(45.42 \pm 8.67)$ .

**Table1. Correlation between neck pain and hand grip strength.**

Grip strength	Neck pain	p-value
$33.42 \pm 13.05$	$3.58 \pm 1.88$	0.0001




Figure 3. Correlation between neck pain and hand grip strength.

Table 2. comparison between mean value of pain intensity in different neck position.

	Flexion	Side bending	Rotation with flexion
Pain intensity Mean±SD	3.55±1.7	3.14±1.5	4.33±3.3
	<i>p</i> -value	0.662	




Fig4. comparison between mean value of pain intensity in different neck position.

Table 3. comparison between mean value of hand grip strength in different neck position

	Flexion	Side bending	Rotation with flexion
Hand grip strength Mean±SD	31.35±12.04	38.28±15.56	45.42±8.67
	<i>p</i> -value	0.197	




Figure 5 comparison between mean value of hand grip strength in different neck position

#### 4 DISCUSSION

In this study we examined the relationship between neck pain intensity and the hand grip strength in dentists. The results revealed that there was a significant positive correlation between neck pain and hand grip strength. As increase in intensity of neck pain lead to increase grip strength. This findings may be attributed to abnormality of sensory motor integration found in neck pain lead to deficit in the quality of sensory information that generate motor output. Also, it can be attributed to sensory hyper excitability. This result agreed with (MaaikA. etal)<sup>28</sup>. They measured the grip strength in neck pain subjects and control group while carrying a 300g weight connected to dynamometer which measured forces applied to the object. They found that subjects with neck pain used significantly higher grip forces as compared to subjects without pain. After the initial lift object, all subject groups adapted to the lifting task by lowering their grip force level but grip force levels were still significantly higher in the subjects with pain.

Our results disagree with (Michael. O. etal)<sup>(29)</sup> who found that the grip strength decrease significantly in patients with spondylosis compared with those control group in dominant and non-dominant hand.

They clarified that increased tissue pressure arising from degenerative changes compromises myoneural conduction velocity and tissue blood flow and oxygenation. These factors interfere with the ability of the nervous system to activate hand muscles.

The results of our study showed that most pain intensity was found in rotation combined with flexion position compared with flexion and side bending positions taken by dentists. These results may be due to strain on ipsilateral facet joint which increase the risk of facet joint injury found in neck rotation.<sup>(30, 31)</sup> Other study demonstrated that at 20° and 40° of rotation of the head decrease the ipsilateral neuro foraminal size up to 15% and 23%, compared with the neutral position.<sup>(32)</sup> While in neck flexion they found that increase in posterior tensile force on spinal canal and soft tissue structure<sup>(33)</sup>. The combination of two movements make much load on the cervical structure which cause more pain.

#### 5 CONCLUSION

It was concluded that there was a significant direct proportional correlation between neck pain and hand grip strength in dentists.

#### RECOMMENDATIONS

It was recommended that the same study have to be conducted on larger sample size, and on different degrees of neck positions.

We recommended the dentists that avoid taking same position during work for long time and take rest in between to prevent the consequences on their hand grip strength during different dental procedures.

**ACKNOWLEDGMENT**

I would like to acknowledge Anwar Adnan M al mousa, FatimahAlwi M Al sadah,Toka Al Awazim, Noor Haddad, Noor Hassan Albrahim, Nawal Al shemary. For their great help during this research project.We also want to thank dentists how participant in our project and help us to collect data.

**REFERENCES**

- [1] Fejer R. Neck pain (Prevalence, genetic and environmental factors). *Ugeskrift for læger*, vol.168, pp.3232, 2013.
- [2] Johnson H, Linda C. The Burden and Determinants of Neck Pain in the General Population. *European Spine Journal*,vol.17,pp.39-51, 2008.
- [3] Jensen I. Neck pain. *Best Practice & Research Clinical Rheumatology*, vol.21,pp. 93-108, 2007.
- [4] Fejer R, Kyvik KO, Hartvigsen J. The prevalence of neck pain in the world population: a systematic critical review of the literature. *European spine journal*, vol.15,pp.834-48, 2006.
- [5] Bernard M, Karnath M. Identifying the Musculoskeletal Causes of Neck Pain. *the Journal of musculoskeletal medicine*,vol.29,pp.82-6,2012.
- [6] Hush JM, Michaleff Z, Maher CG, Refshauge K. Individual, physical and psychological risk factors for neck pain in Australian office workers: a 1-year longitudinal study. *European spine journal*, vol.18, pp.1532-40, 2009.
- [7] Morse T, Bruneau H, Dusssetschleger J. Musculoskeletal disorders of the neck and shoulder in the dental professions. *Occupational and Environmental Health Center*,vol. 35,pp. 419–29, 2010.
- [8] Al Wazzan K, Almas K, AlShethri S, Al Qahtani M. Back & Neck Problems Among Dentists and Dental Auxiliaries. *The Journal of Contemporary Dental Practice*,vol 1,pp.2:10,2001.
- [9] Külçü D, Güluen G, Altunok T, Küçükoulu D, Naderi S. Neck and Low Back Pain Among Dentistry Staff. *Turk J Rheumatol*,vol.25,pp. 9-122, 2010.
- [10] Finsen L, Christensen H, Bakke M. Musculoskeletal disorders among dentists and variation in dental work. *National Institute of Occupational Health, Department of Physiology Lersø Parkalle 105, Copenhagen DK-2100, Denmark*,vol. 29,pp.119-25, 1998.
- [11] Shrestha B, Singh G, Niraula S. Work Related Complaints among Dentists. *J Nepal Med Assoc*,vol.47,pp. 77-81, 2008.
- [12] Barut C, Demirel P. Influence of testing posture and elbow position on grip strength. *Medical Journal of Islamic World Academy of Sciences*,vol,20,pp. 94-7, 2012.
- [13] Massy-Westropp N, Gill T, Taylor A, Bohannon R, Hill C. Hand Grip Strength: age and gender stratified normative data in a population-based study. *BMC Research Notes*, vol. 4,pp.127:35,2011.
- [14] Hanten W, Chen W, Austin A, Brooks R, Carter H, Law C, et al. Maximum Grip Strength in Normal Subjects from 20 to 64 Years of Age *J HAND THER*,vol.12,pp.193-200, 1999.
- [15] Demura S, Aoki H, Sugiura H. Gender differences in hand grip power in the elderly. *Archives of Gerontology and Geriatrics*,vol.53,pp.76-8, 2011.
- [16] Bohannon R, Peolsson A, Westropp N, Desrosiers J, Lehman J. Reference values for adult grip strength measured with a Jamar dynamometer: a descriptive meta-analysis. *Physiotherapy journal*, Vol.92, pp.11–5, 2006.
- [17] Adedoyin R, Ogundapo F, Mbada C, Adekanla B, Olubusola J, Onigbinde T, et al. Reference values for hand grip strength among healthy adults in Nigeria. *Hong Kong Physiother J*,vol.27,pp. 21-9, 2009
- [18] Oxford K. Elbow Positioning for Maximum Grip Performance. *Journal of hand therapy*, vol.13,pp.33-6, 2000.
- [19] Langhammer B, Lindmark B, Stanghelle J. Stroke patients and long-term training is it worthwhile? A randomized comparison of two different training strategies after rehabilitation. *clinical rehabilitation*,vol.21,pp. 495–510, 2007.
- [20] Bohannon R. is it legitimate to characterize muscle strength using a limited number of measures?. *Journal of Strength and Conditioning Research*,vol.22,pp166-73, 2008.
- [21] Wilgen C, Akkerman L, Wieringa J, Dijkstra P. Muscle strength in patients with chronic pain. *Clinical Rehabilitation*,vol.17,pp 885–9, 2003.
- [22] Seo NJ, Sindhu BS, Shechtman O. Influence of pain associated with musculoskeletal disorders on grip force timing. *J HAND THER*,vol.24,pp.335-43,2011.
- [23] Kanea R, Bershadska B, Rockwooda T, Saleha K, Islama N. Visual Analog Scale pain reporting was standardized. *Journal of clinical epidemiology*,vol.58,pp.618–23. 2005.
- [24] Clair D, Edmondston S, Allison G. Variability in pain intensity, physical and psychological function in non-acute, non-traumatic neck pain. *Physiotherapy Research International*,vol.9,pp.43–54,2004.
- [25] Huysmans MA, Hoozemans MJM, Visser B, van Dieën JH. Grip force control in patients with neck and upper extremity pain and healthy controls. *Clinical Neurophysiology*.vol.119,pp.1840-8, 2008.

- [26] Egwu MO, Ajao BA, Mbada CE, Adeosun IO. Isometric Grip Strength and Endurance of Patients With Cervical Spondylosis and Healthy Controls: A Comparative Study. *Hong Kong Physiotherapy Journal*,vol.27,pp.2-6,2009.
- [27] van Eerd M, Patijn J, Lataster A, Rosenquist RW. Cervical facet pain. *Pain practice*,vol.10,pp.113-23, 2010.
- [28] Storvik SG, Stemper BD. Axial head rotation increases facet joint capsular ligament strains in automotive rear impact. *Medical and Biological Engineering and Computing*,vol.49,pp.153-61, 2011.
- [29] Muhle C, Resnick D, Ahn JM, Südmeyer M. In Vivo Changes in the Neuroforaminal Size at Flexion-Extension and Axial Rotation Of The Cervical Spine In Healthy Persons Examined Using Kinematic Magnetic Resonance Imaging. *Spine*, vol.26,pp.e287-e93, 2001.
- [30] Marshall KG. THE CERVICAL SPINE. *Patient Care*,vol.14,pp.S1, 2003.